

Treating ADHD With Suggestion: Neurofeedback and Placebo Therapeutics

Journal of Attention Disorders
2018, Vol. 22(8) 707–711
© The Author(s) 2018
Reprints and permissions:
sagepub.com/journalsPermissions.nav
DOI: 10.1177/1087054718770012
journals.sagepub.com/home/jad



Robert T. Thibault^{1,2} , Samuel Veissière¹,
Jay A. Olson¹, and Amir Raz^{1,2}

Abstract

Objective: We propose that clinicians can use suggestion to help treat conditions such as ADHD. **Methods:** We use EEG neurofeedback as a case study, alongside evidence from a recent pilot experiment utilizing a sham MRI scanner to highlight the therapeutic potential of suggestion-based treatments. **Results:** The medical literature demonstrates that many practitioners already prescribe treatments that hardly outperform placebo comparators. Moreover, the sham MRI experiment showed that, even with full disclosure of the procedure, suggestion alone can reduce the symptomatology of ADHD. **Conclusion:** Non-deceptive suggestion-based treatments, especially those drawing on accessories from neuroscience, may offer a safe complement and potential alternative to current standard of care for individuals with ADHD. (*J. of Att. Dis.* 2018; 22(8) 707-711)

Keywords

placebo, nocebo, neuroenchantment, neurosuggestion, neurofeedback, ADHD, EEG

Recent critical publications on neurofeedback raise a conundrum: Does electroencephalography-neurofeedback (EEG-nf) work, and if so, how? In a series of papers, we reported that EEG-nf seems to help the symptoms of ADHD, but for different reasons than most advocates would expect—that is, due to a placebo response (Thibault, Lifshitz, Birbaumer, & Raz, 2015; Thibault & Raz, 2017). In this Guest Editorial, we address ethical considerations concerning prescribing placebos and highlight how we can leverage prevailing brain-based beliefs about behavioral disorders to better treat patients. We conclude that clinicians can apply EEG-nf to ADHD as a form of clinical suggestion for patients with the time, finances, and inclination to pursue such a treatment.

In a therapeutic context, suggestion refers to communicable ideas, contextual cues, and cultural rituals that can help modulate both voluntary actions and autonomic functions that typically lie outside of our volitional control. Suggestions come in many flavors (e.g., hypnotic, parental, or commercial). In this paper, we focus on the kind of suggestion that uses ideas, cues, and rituals drawn from neurotechnology and our culturally widespread trust in brain-based explanatory models of human behavior—that is, *neurosuggestion*. To enhance the involvement of the patient and to make suggestion more effective, we can leverage culturally salient props, such as a brain scanner, in a process we broadly term *accessory-assisted healing*.

Why EEG-nf Is Neurosuggestion Therapy

EEG-nf for ADHD shows comparable benefits whether the feedback is genuine (from one's own brain signal of interest) or a sham (from an unrelated signal). To date, every relevant double-blind sham-controlled study has reached this conclusion (Arnold et al., 2013; Lansbergen, van Dongen-Boomsma, Buitelaar, & Slaats-Willemse, 2011; Logemann, Lansbergen, Van Os, Bocker, & Kenemans, 2010; Perreau-Linck, Lessard, Lévesque, & Beaugard, 2010; Schönenberg et al., 2017; Thibault & Raz, 2017; van Dongen-Boomsma, Vollebregt, Slaats-Willemse, & Buitelaar, 2013; Vollebregt, van Dongen-Boomsma, Buitelaar, & Slaats-Willemse, 2014). In all but two of these studies, the treatment benefited both groups (Logemann et al., 2010; Vollebregt et al., 2014). A recent triple-blind registered randomized controlled trial (RCT) with 118 participants diagnosed with ADHD found large effect sizes for both true ($d = 1.0$) and sham ($d = 1.5$) neurofeedback groups (Schönenberg et al., 2017). Until research demonstrates additional benefits from true neurofeedback that go

¹McGill University, Montreal, Québec, Canada

²Chapman University, Irvine, CA, USA

Corresponding Author:

Robert T. Thibault, Brain Institute, Chapman University, Irvine, CA 92618, USA.

Email: robert.thibault@mail.mcgill.ca

above and beyond placebo effects, the science suggests that EEG-nf, at least in its current incarnation, relies exclusively on treatment mechanisms unrelated to watching one's own brain activity (Thibault & Raz, 2016). EEG-nf, in other words, is neurosuggestion.

We Already Prescribe Suggestion

Most clinicians have prescribed suggestion—but many don't know it. The benefits of selective serotonin reuptake inhibitors (SSRIs) for depression, acupuncture for lower back pain, and knee surgery for osteoarthritis all stem largely from the suggestion that these treatments will improve your health (i.e., from placebo effects). In 2002, the psychologist Irving Kirsch obtained unpublished clinical trial data under the Freedom of Information Act and found that SSRIs barely outperformed placebos in RCTs (Kirsch, Moore, Scoboria, & Nicholls, 2002). In the case of lower back pain, two RCTs with a total of more than 1,700 participants demonstrated comparable benefits between veridical acupuncture and sham acupuncture, yet both outperformed a standard-of-care treatment (Cherkin et al., 2009; Haake et al., 2007; Harris, Lifshitz, & Raz, 2015). Likewise, a high-profile RCT showed that, even at a 2-year follow-up, sham knee surgery decreased pain and improved movement capacity on par with real surgery (Moseley et al., 2002). An expert panel now strongly recommends against knee surgery for osteoarthritis due to potential complications and the absence of benefits beyond placebo effects (Siemieniuk et al., 2017). In these situations, clinicians often unknowingly prescribe placebos. Following the results from recent double-blind studies, we can now add EEG-nf for ADHD to this list of placebo therapies that masquerade under other biomedical labels.

The Case for Suggestion in ADHD

In contrast to the above-mentioned placebo therapies, psychostimulants typically reduce ADHD symptoms more effectively than placebo, but come at a cost. Potential side effects include crying, staring, anxiety, sadness, nail biting, euphoria, and shyness (Konrad-Bindl, Gresser, & Richartz, 2016). Due to growing concerns about long-term adverse side effects, the European Commission recently called for a 2-year longitudinal study that is taking place at 27 sites (Inglis et al., 2016). Given the potential for harm, it is advantageous to consider drug-free treatments with minimal side effects to complement or replace psychostimulants. In this vein, a pair of studies found that when children with ADHD were prescribed an "open-label" placebo, they could cut their psychostimulant intake in half with negligible changes in behavior (Sandler & Bodfish, 2008; Sandler, Glesne, & Bodfish, 2010). In lieu of deception, the researchers briefed children with the following script:

This little capsule is a placebo. Placebos have been used a lot in treating people. It is called "Dose Extender." As you can see, it is different from Adderall. Dose Extender is something new. It has no drug in it. I can promise you that it won't hurt you at all. It has no real side effects. But it may help you to help yourself. It may work well with your Adderall, kind of like a booster to the dose of Adderall. That's why it's called a Dose Extender. I won't be surprised when I hear from you and your parents and your teachers that you're able to control your ADHD better. (Sandler & Bodfish, 2008, p. 106)

Full disclosure made little impact on the effectiveness of the placebo condition—that is to say, administering placebos openly hardly detracted from the clinical benefit (Sandler & Bodfish, 2008). Researchers have reproduced this finding in several conditions, including irritable bowel syndrome, chronic pain, and depression (Kaptchuk et al., 2010; Kelley, Kaptchuk, Cusin, Lipkin, & Fava, 2012; Schafer, Colloca, & Wager, 2015).

Based on staunch faith in brain science, neurosuggestion may perhaps treat patients even better than placebo pills. In particular, with EEG-nf for ADHD, the suggestion that physical movement will contaminate the expensive brain recordings can lead participants to sit still and in turn, this procedure provides an ulterior form of behavioral therapy. The additional psychosocial cues surrounding EEG-nf, compared with those present when ingesting a pill, may help this technique compete with standard pharmacotherapy (e.g., Fuchs, Birbaumer, Lutzenberger, Gruzelier, & Kaiser, 2003). To further support this point scientifically and to the level of clinical recommendation, more studies would need to directly compare EEG-nf with standard of care treatment.

Testing Neurosuggestion in the Clinic

In our previous work, we explored how strong this type of suggestion could be. We found that many participants believed that a crudely built "brain scanner"—assembled from discarded hair drying and medical equipment and placed in a cognitive neuroscience laboratory—could read their minds (Ali, Lifshitz, & Raz, 2014). In subsequent iterations of this paradigm, we found that a more realistic looking scanner could similarly alter people's sense of agency and moral attitudes (Olson, Landry, Appourchaux, & Raz, 2016; Olson, Strandberg, Hall, Johansson, & Raz, 2017). In light of these findings, we decided to test the clinical applications of this technique we term neurosuggestion.

To pilot the idea, we used a real but decommissioned—in other words, sham—magnetic resonance imaging (MRI) scanner at the Montreal Neurological Institute as part of an open-label procedure to treat nine children diagnosed with ADHD (Figure 1; Veissière, Olson, & Raz, 2017). We told the children that the "brain machine" was inactive, and that we would "use it as a suggestion" that would "help their



Figure 1. Decommissioned Siemens 1.5T MRI used for the neurosuggestion procedure.

Note. MRI = magnetic resonance imaging.

brain heal itself.” While in the defunct scanner, we gave the children positive verbal suggestions to promote relaxation, focus, and confidence. At 1-, 3-, and 6-week follow-ups, parents reported improvements in eight out of the nine participating children. In qualitative interviews, two families reported near complete remission of symptoms, and six reported improvements in areas such as confidence, self-control, and social skills. None reported any side effects. In essence, this study provided neurofeedback-like treatment, but instead of focusing on a specific physiological mechanism, we emphasized suggestion-based healing.

Should We Prescribe EEG-nf for ADHD?

To answer this question, we need to consider a few issues. First, as a culture, we maintain deep-set beliefs that the origin of behavioral disorders resides in the brain (Moncrieff, 2016; Rose, 2003). Relying on this assumption, children and their families often actively seek a diagnostic label, for example, ADHD, to ascribe meaning to their behavior (Moncrieff, 2016; Moncrieff, Rapley, & Timimi, 2015). Second, even in our open-label neurosuggestion experiment (Veissière et al., 2017), parents continued to ask what was “wrong” with their child’s brain, despite having been extensively briefed regarding the inert nature of the MRI scanner. And yet, this very belief system—that a brain disorder is the core reason for the symptoms—may unintentionally act as a suggestion to further obfuscate the situation and exacerbate symptoms through nocebo effects (Loftus & Fries, 1979). This type of thinking has been described as a “looping effect,” where our beliefs and social norms affect the framing, course, and outcomes of

a disorder (Hacking, 1995). In other words, context and culture strongly modulate how patients attend to and construe their behavior, which, in turn, alters their symptoms (Kirmayer, Gomez-Carrillo, & Veissière, 2017; Seth & Friston, 2016). Third, our experiments show that brain-based folk explanations can lead to both nocebo and placebo effects. It seems that neurofeedback likely summons its strength from this belief system, which offers a promising means to regulate “faulty” brain patterns (Thibault & Raz, 2017). Thus, neurosuggestion, rather than the act of regulating one’s own EEG waves, likely allows patients to break free from harmful looping effects.

In sum, under certain circumstances, clinicians could ethically and nondeceptively prescribe EEG-nf as a form of neurosuggestion therapy. It would not be the first placebo prescription—in the United States, one study reports that 45% of physicians use placebos in clinical practice and 96% of them believe placebos can have therapeutic effects (Sherman & Hickner, 2008); in Canada, an analysis of placebo use among physicians, and especially psychiatrists, echoes similar sentiments (Raz et al., 2011). Moreover, a recent study describes interviews with 1,000 parents and reports that most deem it acceptable for clinicians to recommend placebo treatment for ADHD (Faria et al., 2017). With these realities in mind, EEG-nf presents a reasonable alternative for patients experiencing adverse side effects or simply searching for a nonpharmacological treatment. Because EEG-nf requires time, money, and energy, if clinicians decide to promote this technique, they ought to present it as one of several options (e.g., exercise, cognitive training, behavioral therapy, and diet; Sonuga-Barke et al., 2013). Thus, clinicians can certainly prescribe EEG-nf. We recommend they do so transparently and with an eye for amplifying the psychosocial mechanisms of suggestion rather than grasping at the elusive neural signatures many practitioners speciously assign as the cause of ADHD.

Author Contribution

Robert T. Thibault and Samuel Veissière contributed equally to this work.

Declaration of Conflicting Interests


The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: RTT acknowledges an Alexander Graham Bell Canada Graduate Scholarship and Micheal Smith Foreign Study Supplement from the Natural Sciences and Engineering Research Council of Canada (NSERC). SV and JAO received funding and logistical support from the Mind Field segment of the Intellectual Property Corporation for the sham MRI experiment, which is approved under McGill

University REB File: #32-0617. JAO acknowledges funding from the Joseph-Armand Bombardier Scholarship from the Social Sciences and Humanities Research Council of Canada. We further thank the BIAL Foundation for funding. The funding sources had no involvement in reviewing the literature, writing the manuscript, or deciding to submit the paper for publication.

ORCID iD

Robert T. Thibault  <https://orcid.org/0000-0002-6561-3962>

References

- Ali, S., Lifshitz, M., & Raz, A. (2014). Empirical neuroenchantment: From reading minds to thinking critically. *Frontiers in Human Neuroscience*, *8*, Article 357. doi:10.3389/fnhum.2014.00357
- Arnold, L. E., Lofthouse, N., Hersch, S., Pan, X., Hurt, E., Bates, B., . . . Grantier, C. (2013). EEG neurofeedback for ADHD: Double-blind sham-controlled randomized pilot feasibility trial. *Journal of Attention Disorders*, *17*, 410-419. doi:10.1177/1087054712446173
- Cherkin, D. C., Sherman, K., Avins, A., Erro, J., Ichickawa, L., . . . Deyo, R. A. (2009). A randomized trial comparing acupuncture, simulated acupuncture, and usual care for chronic low back pain. *Archives of Internal Medicine*, *169*, 858-866. doi:10.1001/archinternmed.2009.65
- Faria, V., Kossowsky, J., Petkov, M. P., Kaptchuk, T. J., Kirsch, I., Lebel, A., & Borsook, D. (2017). Parental attitudes about placebo use in children. *Journal of Pediatrics*, *181*, 272-278. doi:10.1016/j.jpeds.2016.10.018
- Fuchs, T., Birbaumer, N., Lutzenberger, W., Gruzelier, J. H., & Kaiser, J. (2003). Neurofeedback treatment for attention-deficit/hyperactivity disorder in children: A comparison with methylphenidate. *Applied Psychophysiology and Biofeedback*, *28*, 1-12.
- Haake, M., Müller, H., Schade-Brittinger, C., Basler, H., Schäfer, H., Maier, C., . . . Molsberger, A. (2007). German Acupuncture Trials (GERAC) for chronic low back pain: randomized, multicenter, blinded, parallel-group trial with 3 groups. *Archives of Internal Medicine*, *167*, 1892-1898. doi:10.1001/archinte.167.17.1892
- Hacking, I. (1995). The looping effects of human kinds. In Sperber D., Premack D., & Premack A. J. (Eds.) *Causal cognition: A Multidisciplinary Debate* (pp. 351-394). New York, NY: Clarendon Press/Oxford University Press.
- Harris, C. S., Lifshitz, M., & Raz, A. (2015). Acupuncture for chronic pain? Clinical wisdom undecided despite over 4000 years of practice. *The American Journal of Medicine*, *128*, 331-333. doi:10.1016/j.amjmed.2014.10.042
- Inglis, S. K., Carucci, S., Garas, P., Häge, A., Banaschewski, T., Buitelaar, J. K., . . . Coghill, D. C. (2016). Prospective observational study protocol to investigate long-term adverse effects of methylphenidate in children and adolescents with ADHD: The Attention Deficit Hyperactivity Disorder Drugs Use Chronic Effects (ADDUCE) study. *BMJ Open*, *6*(4), 1-12. doi:10.1136/bmjopen-2015-010433
- Kaptchuk, T. J., Friedlander, E., Kelley, J. M., Sanchez, M. N., Kokkotou, E., Singer, J. P., . . . Lembo, A. J. (2010). Placebos without deception: A randomized controlled trial in irritable bowel syndrome. *PLoS ONE*, *5*(12), Article e15591. doi:10.1371/journal.pone.0015591
- Kelley, J. M., Kaptchuk, T. J., Cusin, C., Lipkin, S., & Fava, M. (2012). Open-label placebo for major depressive disorder: A pilot randomized controlled trial. *Psychotherapy and Psychosomatics*, *81*, 312-314. doi:10.1159/000337053
- Kirmayer, L. J., Gomez-Carrillo, A., & Veissière, S. (2017). Culture and depression in global mental health: An ecosocial approach to the phenomenology of psychiatric disorders. *Social Science & Medicine*, *183*, 163-168. doi:10.1016/j.socscimed.2017.04.034
- Kirsch, I., Moore, T. J., Scoboria, A., & Nicholls, S. S. (2002). The emperor's new drugs: An analysis of antidepressant medication data submitted to the U.S. food and drug administration. *Prevention & Treatment*, *5*(1), 1-11. doi:10.1037//1522-3736.5.1.523a
- Konrad-Bindl, D. S., Gresser, U., & Richartz, B. M. (2016). Changes in behavior as side effects in methylphenidate treatment: Review of the literature. *Neuropsychiatric Disease and Treatment*, *12*, 2635-2647. doi:10.2147/NDT.S114185
- Lansbergen, M. M., van Dongen-Boomsma, M., Buitelaar, J. K., & Slaats-Willemse, D. (2011). ADHD and EEG-neurofeedback: A double-blind randomized placebo-controlled feasibility study. *Journal of Neural Transmission*, *118*, 275-284. doi:10.1007/s00702-010-0524-2
- Loftus, E. F., & Fries, J. F. (1979). Informed consent may be hazardous to health. *Science*, *204*(4388), 11. doi:10.1126/science.373117
- Logemann, H. N. A., Lansbergen, M. M., Van Os, T. W. D. P., Bocker, K. B. E., & Kenemans, J. L. (2010). The effectiveness of EEG-feedback on attention, impulsivity and EEG: A sham feedback controlled study. *Neuroscience Letters*, *479*, 49-53. Retrieved from <http://www.sciencedirect.com/science/article/pii/S0304394010006014>
- Moncrieff, J. (2016). *The myth of the chemical cure: A critique of psychiatric drug treatment*. Basingstoke, UK: Pgrave Macmillan.
- Moncrieff, J., Rapley, M., & Timimi, S. (2015). The construction of psychiatric diagnoses: The case of adult ADHD*. *The Journal of Critical Psychology, Counselling and Psychotherapy*, *15*, 42-55.
- Moseley, J. B., O'Malley, K., Petersen, N. J., Menke, T. J., Brody, B. A., Kuykendall, D. H., . . . Wray, N. P. (2002). A controlled trial of arthroscopic surgery for osteoarthritis of the knee. *The New England Journal of Medicine*, *347*, 81-88. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/12368798>
- Olson, J. A., Landry, M., Appourchoux, K., & Raz, A. (2016). Simulated thought insertion: Influencing the sense of agency using deception and magic. *Consciousness and Cognition*, *43*, 11-26. doi:10.1016/j.concog.2016.04.010
- Olson, J. A., Strandberg, T., Hall, L., Johansson, P., & Raz, A. (2017, October). *Manipulating attitudes using suggestion*. 68th Society for Clinical and Experimental Hypnosis Conference, Chicago, IL.
- Perreau-Linck, E., Lessard, N., Lévesque, J., & Beauregard, M. (2010). Effects of neurofeedback training on inhibitory capacities in ADHD children: A single-blind, randomized,

- placebo-controlled study. *Journal of Neurotherapy*, *14*, 229-242. doi:10.1080/10874208.2010.501514
- Raz, A., Campbell, N., Guindi, D., Holcroft, C., Déry, C., & Cukier, O. (2011). Placebos in clinical practice: Comparing attitudes, beliefs, and patterns of use between academic psychiatrists and nonpsychiatrists. *Canadian Journal of Psychiatry*, *56*, 198-208. doi:10.1177/070674371105600403
- Rose, N. (2003). Neurochemical selves. *Society*, *41*, 46-59. doi:10.1007/BF02688204
- Sandler, A., & Bodfish, J. W. (2008). Open-label use of placebos in the treatment of ADHD: A pilot study. *Child: Care, Health and Development*, *34*, 104-110. doi:10.1111/j.1365-2214.2007.00797.x
- Sandler, A., Glesne, C. E., & Bodfish, J. W. (2010). Conditioned placebo dose reduction: A new treatment in attention-deficit hyperactivity disorder? *Journal of Developmental and Behavioral Pediatrics*, *31*, 369-375. doi:10.1097/DBP.0b013e3181e121ed
- Schafer, S. M., Colloca, L., & Wager, T. D. (2015). Conditioned placebo analgesia persists when subjects know they are receiving a placebo. *Journal of Pain*, *16*, 412-420. doi:10.1016/j.jpain.2014.12.008
- Schönenberg, M., Wiedemann, E., Schneidt, A., Scheeff, J., Logemann, A., Keune, P. M., & Hautzinger, M. (2017). Neurofeedback, sham neurofeedback, and cognitive-behavioural group therapy in adults with attention-deficit hyperactivity disorder: A triple-blind, randomised, controlled trial. *The Lancet Psychiatry*, *4*, 673-684.
- Seth, A. K., & Friston, K. J. (2016). Active interoceptive inference and the emotional brain. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *371*(1708), Article 20160007. doi:10.1098/rstb.2016.0007
- Sherman, R., & Hickner, J. (2008). Academic physicians use placebos in clinical practice and believe in the mind-body connection. *Journal of General Internal Medicine*, *23*, 7-10. doi:10.1007/s11606-007-0332-z
- Siemieniuk, R. A. C., Harris, I. A., Agoritsas, T., Poolman, R. W., Brignardello-Petersen, R., Van de Velde, S., . . . Kristiansen, A. (2017). Arthroscopic surgery for degenerative knee arthritis and meniscal tears: A clinical practice guideline. *British Medical Journal*, *357*, Article j1982. doi:10.1136/BMJ.J1982
- Sonuga-Barke, E. J. S., Brandeis, D., Cortese, S., Daley, D., Ferrin, M., Holtmann, M., . . . Sergeant, J. (2013). Nonpharmacological interventions for ADHD: Systematic review and meta-analyses of randomized controlled trials of dietary and psychological treatments. *The American Journal of Psychiatry*, *170*, 275-289. doi:10.1176/appi.ajp.2012.12070991
- Thibault, R. T., Lifshitz, M., Birbaumer, N., & Raz, A. (2015). Neurofeedback, self-regulation, and brain imaging: Clinical science and fad in the service of mental disorders. *Psychotherapy and Psychosomatics*, *84*, 193-207. doi:10.1159/000371714
- Thibault, R. T., & Raz, A. (2016). Neurofeedback: The power of psychosocial therapeutics. *The Lancet Psychiatry*, *3*(11), Article e18.
- Thibault, R. T., & Raz, A. (2017). The psychology of neurofeedback: Clinical intervention even if applied placebo. *American Psychologist*, *72*, 679-688. doi:10.1037/amp0000118
- van Dongen-Boomsma, M., Vollebregt, M. A., Slaats-Willemse, D., & Buitelaar, J. K. (2013). A randomized placebo-controlled trial of Electroencephalographic (EEG) neurofeedback in children with attention-deficit/hyperactivity disorder. *Journal of Clinical Psychiatry*, *74*, 821-827. doi:10.4088/JCP.12m08321
- Veissière, S., Olson, J. A., & Raz, A. (2017, October). *Neurosuggestion improves self-regulation in neurodevelopmental disorders: A feasibility study*. 68th annual meeting of the Society for Clinical and Experimental Hypnosis, Chicago, IL.
- Vollebregt, M. A., van Dongen-Boomsma, M., Buitelaar, J. K., & Slaats-Willemse, D. (2014). Does EEG-neurofeedback improve neurocognitive functioning in children with attention-deficit/hyperactivity disorder? A systematic review and a double-blind placebo-controlled study. *Journal of Child Psychology and Psychiatry and Allied Disciplines*, *55*, 460-472. doi:10.1111/jcpp.12143

Author Biographies

Robert T. Thibault is a PhD Candidate in the Integrated Program in Neuroscience at McGill University. He is interested in expediting scientific discoveries and improving medical treatment by leveraging biases and heuristics when helpful and circumventing them when harmful.

Samuel Veissière is an anthropologist and cognitive scientist by training and currently an Assistant Professor in the Culture, Mind, and Brain program in the Department of Psychiatry at McGill University. He studies how culture modulates cognition, attention, and healing from evolutionary and experimental perspectives.

Jay A. Olson is a PhD candidate in Psychiatry at McGill University. He studies how to maximize the effects of suggestion and placebos.

Amir Raz is a Professor in the Department(s) of Psychiatry (Neurology & Neurosurgery, and Psychology) at McGill University; Senior Researcher at the Lady Davis Institute for Medical Research and the Institute for Community and Family Psychiatry at the Jewish General Hospital in Montreal, Quebec, Canada; and Director and Professor of the Brain Institute at Chapman University, Irvine, CA, USA.